



Early accretion of planetesimals unraveled by the thermal evolution of the parent bodies of magmatic iron meteorites

Edouard Kaminski^{*}, Angela Limare, Balthasar Kenda, Marc Chaussidon

Université de Paris, Institut de Physique du Globe de Paris, CNRS, 1 rue Jussieu, F-75005 Paris, France

ARTICLE INFO

Article history:

Received 20 February 2019

Received in revised form 7 July 2020

Accepted 8 July 2020

Available online 31 July 2020

Editor: W.B. McKinnon

Keywords:

planetesimals

accretion

convection

analog experiments

magmatic iron meteorites

ABSTRACT

The timing of formation of 100–300 km size planetesimals in the protoplanetary disk remains largely unconstrained. Recent models show that gravitational collapse of boulders in overdense regions of a dusty accretion disk can overcome the meter-sized barrier and lead to rapid formation of planetesimals with size of several km that further grow by pebble accretion. Hf/W ages indicate that the first large planetesimals to form could be the parent bodies of magmatic iron meteorites. These ages have been so far used to constrain timing of accretion considering (i) instantaneous accretion, and (ii) purely conductive heat transfer in the planetesimal. To relax these hypotheses we model the thermal evolution of a planetesimal in course of accretion and we take into account the possibility of convection onset. Our model is further based on considering the possibility of a common thermal evolution for all the parent bodies of iron meteorites. Within that framework we show that the planetesimals could have grown following a universal accretion law starting at the very beginning of the history of the disk by a nearly instantaneous formation of 60 ± 30 km size nuclei, followed by a growth via pebble accretion at a much slower pace to reach final sizes of 150–300 km in about 3 Myr. In this universal scenario, complete melting and total differentiation are not bound to happen in the parent body due to the continuous accretion of cold pebbles. The model, though calibrated here on iron meteorites, is general and can in principle be applied to other types of planetesimals such as for instance the parent bodies of CV chondrites.

© 2020 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The formation of planetesimals is a crucial stage in the evolution of the protoplanetary disk that leads from dust to planets. Some of its fundamental aspects remain the object of debate, such as the range in initial and final sizes of the planetesimals, and more generally their mechanism and rate of accretion (e.g., Xie et al., 2010; Morbidelli and Raymond, 2016; Birnstiel et al., 2016).

A key and long-standing question has been the process of formation of the initial seeds of the planetesimals. Models of collision between aggregates of dust held by electrostatic forces fail to produce planetesimals, because the particles start to bounce upon collision when they reach few centimeters and migrate rapidly toward the star as they grow (e.g., Morbidelli and Raymond, 2016). Meter-size boulders migrate so fast that they are eventually lost to the sun. One way to overcome this meter-size barrier (Weidenschilling, 1977) is to consider a weakly turbulent nebula where large and dense clumps of chondrules can form on the smallest scale of the

flow, and be preserved by self-gravity to later produce planetesimals by sedimentation towards their center (Cuzzi et al., 2008). Numerical simulations (e.g., Johansen et al., 2015; Simon et al., 2017) show that other mechanisms such as quasi-stable pressure bumps and/or streaming instabilities can produce gravitationally-bound pebble-swarms collapse under their own gravity, yielding initial planetesimal sizes up to many tens of kilometers across.

Once the seeds of planetesimals are formed, they are supposed to continue to grow by pebble accretion (Johansen and Lambrechts, 2017). This mechanism is controlled by the interplay between gravity, aerodynamics deflection and gas drag (e.g., Ormel and Klahr, 2010). Models predictions give the minimum planetesimal radius for the onset of pebble accretion as well as the growth timescales for a given radius at a given distance from the sun (Visser and Ormel, 2016). But these models have not been used yet to predict complete accretion scenarios for the planetesimals and to confront the output with independent constraints.

Constraints on the accretion of planetesimals in the disk can be obtained by astronomical observations (e.g., van der Marel et al., 2013; Delbo' et al., 2017) as well as by the study of meteoritic fragments that sampled the early stages of the disk evolution dur-

^{*} Corresponding author.

E-mail address: kaminski@ipgp.fr (E. Kaminski).

Table 1

Definition and values of the model parameters or formulae used to compute them. References: ⁺ Sramek et al. (2012), ^o Suzuki et al. (1998), * www.nucleide.org.

Quantity	Symbol	Units	Value/formula
Metal mass ⁺ , volume fraction	$x^{\text{met}}, \phi^{\text{met}}$	–	$x^{\text{met}} = 0.18$
Silicate mass ⁺ , volume fraction	$x^{\text{sil}}, \phi^{\text{sil}}$	–	$x^{\text{sil}} = 0.82$
Average porosity	ϕ	–	0.1
Density of metal ⁺	ρ^{met}	kg m ⁻³	7800
Density of silicate ⁺	ρ^{sil}	kg m ⁻³	3200
Average density	ρ	kg m ⁻³	$1/\rho = x^{\text{met}}/\rho^{\text{met}} + x^{\text{sil}}/\rho^{\text{sil}}$
Specific heat of metal ⁺	C_p^{met}	J K ⁻¹ kg ⁻¹	450
Specific heat of silicate ⁺	C_p^{sil}	J K ⁻¹ kg ⁻¹	1200
Average specific heat	C_p	J K ⁻¹ kg ⁻¹	$C_p = x^{\text{met}}C_p^{\text{met}} + x^{\text{sil}}C_p^{\text{sil}}$
Latent heat of metal ⁺	L^{met}	kJ kg ⁻¹	250
Latent heat of silicate ⁺	C_p^{sil}	kJ kg ⁻¹	500
Activation energy metal	$E_{\text{act}}^{\text{met}}$	kJ mol ⁻¹	13
Activation energy silicate	$E_{\text{act}}^{\text{sil}}$	kJ mol ⁻¹	250
Average activation energy	E_{act}	kJ mol ⁻¹	$E_{\text{act}} = \phi^{\text{met}}E_{\text{act}}^{\text{met}} + \phi^{\text{sil}}E_{\text{act}}^{\text{sil}}$
Thermal conductivity of metal ⁺	k^{met}	J s ⁻¹ m ⁻¹ K ⁻¹	50
Thermal conductivity of silicate ⁺	k^{sil}	J s ⁻¹ m ⁻¹ K ⁻¹	3
Average thermal conductivity	k	J s ⁻¹ m ⁻¹ K ⁻¹	$k = \phi^{\text{met}}k^{\text{met}} + \phi^{\text{sil}}k^{\text{sil}}$
Thermal diffusivity	κ	m ² s ⁻¹	$\kappa = k/(\rho C_p)$
Dynamic viscosity	μ	Pa s	
Kinematic viscosity	ν	m ² s ⁻¹	$\nu = \mu/\rho$
Coefficient of thermal expansion ^o	α	K ⁻¹	$3 \cdot 10^{-5}$
Acceleration of gravity	g	m s ⁻²	$g = 4/3\pi R G \rho$
Universal gravity constant	G	m ³ kg ⁻¹ s ⁻²	$6.674 \cdot 10^{-11}$
Initial radiogenic heat production	H_0	J s ⁻¹ kg ⁻¹	$1.5 \cdot 10^{-7}$
²⁶ Al decay constant*	λ	s ⁻¹	$3 \cdot 10^{-14}$
Radiogenic heat production	H	J s ⁻¹ kg ⁻¹	$H(t) = H_0 \exp(-\lambda t)$
Time since CAI formation	t	s (Myr)	
Parent body average internal temperature	T	°C	
Parent body surface temperature	T_s	°C	0
Parent body surface heat flux	Q_s	J s ⁻¹ m ⁻²	
Parent body radius	R	m	
Initial parent body radius	R_0	m	$R_0 = R(t=0)$
Rayleigh-Roberts number	Ra_H	–	$Ra_H = \frac{\rho g \alpha H R^5}{\kappa \nu}$

ing which planetesimals formed. The very early ages demonstrated by Hf/W studies of iron meteorites presumably originating from planetesimals accreted inside and outside the orbit of the growing Jupiter, indicate that these planetesimals and a Jupiter's core of ≈ 20 Earth mass may have formed within the first million year (Myr) of the solar system (Birnstiel et al., 2016; Kruijer et al., 2014, 2017). At variance, the presence in chondritic meteorites of Ca-, Al-rich inclusions (CAIs) and chondrules spanning in age the first 4 Myr of the solar system (e.g., Villeneuve et al., 2009) shows that their parent planetesimals completed their accretion significantly later. Because no phenomenological law for planetesimal accretion has ever been derived from data on meteorites, the first principles behind these different accretion times still remain inaccessible.

One way to unravel the accretion history of planetesimals is to study their thermal evolution, which is strongly controlled by their size (e.g., Merk et al., 2002) that sets the ratio between internal production of energy (proportional to the cube of the size) and the surface heat loss (proportional to the square of the size). Previous models have essentially considered that the surface heat loss could be estimated based on a conductive profile in the planetesimal (e.g., Ghosh et al., 2003). It has been shown however that melting-induced convection could occur ≈ 1 Myr after the accretion of a planetesimal (Hevey and Sanders, 2006). The conductive hypothesis is thus not valid over the whole thermal history of a planetesimal and more efficient heat transfer by convection must be taken into account. To our knowledge, the recent study of Lichtenberg et al. (2016) is the only one to explicitly model convection in the planetesimal, using 2D and 3D numerical simulations. However this study considers instantaneous accretion and does not tackle the question of the thermal evolution of the planetesimal in the route of accretion. We propose to do so in the present study by

using scaling laws for the transient thermal evolution of an internally heated convective body validated by lab-scale experiments.

2. Thermal evolution of a planetesimal in the course of accretion: principles and constraints

2.1. Conservation of energy

The average internal temperature $T(t)$ of a spherical body of radius $R(t)$ can be defined based on its total thermal energy $E(t)$ as

$$T(t) = \frac{E(t)}{4/3\pi\rho C_p R^3(t)}, \quad (1)$$

with ρ the density and C_p the specific heat (Table 1). The time evolution of this average internal temperature is given by the balance between internal radioactive heating due to decay of short-lived ²⁶Al, $H(t)$, and the surface heat loss, $Q_s(t)$,

$$\rho C_p \frac{4}{3}\pi \frac{dR^3(t)T(t)}{dt} = \frac{4}{3}\pi R^3(t)H(t) - 4\pi R^2(t)Q_s(t) + E_{\text{int}}, \quad (2)$$

where $E_{\text{int}} = E_{\text{lat}} + E_a$ is a term corresponding to the heat absorbed by partial melting (E_{lat}) or brought by surface impacts (E_a), as discussed later (Section 3.1). For a given accretion law $R(t)$, $T(t)$ can be calculated using the energy balance equation (2) if the surface heat flux $Q_s(t)$ is known. Equivalently, but the other way around, if a thermal record $T(t)$ is given, equation (2) can be inverted to obtain the accretion curve $R(t)$.

In the purely conductive regime $Q_s(t)$ is analytically known (Carslaw and Jaeger, 1959, p. 245). In a convective regime, the determination of $Q_s(t)$ requires either the numerical resolution of




Fig. 1. Thermal evolution of the parent body of the iron magmatic meteorites, $T(t) = T_{\infty}[1 - \exp(-t/\tau)]$, with $T_{\infty} = 1524^{\circ}\text{C}$ and $\tau = 0.424$ Myr. The red circles give the position of the “non-carbonaceous” magmatic iron meteorites and the blue circles of the “carbonaceous” (Kruijer et al., 2014; Budde et al., 2016). (For interpretation of the colors in the figure(s), the reader is referred to the web version of this article.)

the full set of conservative equations in the convective fluid (Lichtenberg et al., 2016), or the use of a scaling law relating the surface heat flux to the fluid properties, the internal heating rate, and the geometry of the system; an approach we will detail and follow in section 3.

2.2. Constraints brought by the magmatic iron meteorites

2.2.1. Temperature and age of formation

The ^{182}Hf - ^{182}W age of magmatic iron meteorites can be taken as the formation time of their core, or, more generally, the time of metal-silicate differentiation and metal segregation (Lee and Halliday, 1997; Kleine et al., 2002; Qin et al., 2008; Kruijer et al., 2014). Data for five groups of magmatic iron meteorites (IIAB, IIB, IIIAB, IVA, IVB) provide ages spanning from 0.7 ± 0.3 Myr after CAIs for IIAB to 3.1 ± 0.8 Myr after CAIs for IIB (Kruijer et al., 2014). Their temperatures of formation, taken as the liquidus temperature of the iron-sulfur alloy, range from 1325°C to 1615°C as a function of their sulfur content (see Table 2 from Supplementary material). We note in Fig. 1 that a single exponential function fits within error bars the data on the five groups of magmatic iron meteorites $T(t) = T_{\infty}[1 - \exp(-t/\tau)]$. We used an Orthogonal Distance Regression (ODR) method based on a Levenberg-Marquardt type algorithm that allows non-linear fitting with errors on both variables, and we obtained the following values: $\tau = 0.424 \pm 0.108$ Myr and $T_{\infty} = 1524 \pm 46^{\circ}\text{C}$. Rather than considering a specific thermal evolution hence a somewhat open ended accretion history for each magmatic iron meteorite, such a unique fit opens the possibility to look for a tightly constrained universal accretion scenario that can explain all the data. To that aim, we will use in the following (section 4) the best fitting exponential evolution of the temperature to obtain accretion curves $R(t)$ - if they exist - consistent with this thermal evolution.

2.2.2. Metallographic cooling rates

To further constrain the accretion curves, we will rely on the metallographic cooling rates to infer the size of the parent bodies of magmatic iron meteorites. These correspond to the latest (conductive) cooling phase of a metallic pocket (or core) once it has solidified. They are thus independent from the thermal models considered in this study which will only describe the heating

(and convective) stage of the accreting planetesimals. The size of the planetesimals are poorly constrained and have been estimated based on two assumptions (see Table 3 from Supplementary material).

The IVA meteorites have the highest cooling rates, interpreted as resulting from the conductive cooling of a 150 ± 50 km metallic core with its mantle ripped off, yielding a parent body size of ≈ 300 km (Yang et al., 2008). Cooling rates for IVB meteorites give a parent body size of 140 ± 30 km (Yang et al., 2010).

Other magmatic iron meteorites display smaller cooling rates (Yang and Goldstein, 2006; Goldstein et al., 2009) which reflects the insulating effect of a mantle and possibly of a regolith. Their size depends on the thickness and thermal properties of the rocky envelopes around the core (Haack et al., 1990). Based on the recorded cooling rates and on Fig. 7.1 of McSween (1999) we estimate the radii as 125 ± 50 km for IIIAB, 235 ± 65 km for IIB, and 165 ± 65 km for IIAB.

3. Transient thermal evolution in an internally heated convective system

The resolution of the energy balance equation (2) requires a knowledge of the surface heat flux Q_s which is different in a conductive and in a convective thermal regime. In the case of Rayleigh-Bénard convection, where convection results from cooling at the top of the system and heating at its base, scaling laws are expressed using the Nusselt number, Nu , defined as the ratio between the surface heat flux measured in the convective regime and the surface heat flux in the conductive regime. At steady state, Nu scales as the $1/3$ rd power of the Rayleigh number of the well-mixed convective layer. In internally heated fluids, convection depends on the so called Rayleigh-Roberts number, defined as

$$Ra_H = \frac{\rho g \alpha H R^5}{k \kappa \nu}, \quad (3)$$

with g the gravity, α the coefficient of thermal expansion, k the thermal conductivity, κ the thermal diffusivity and ν the kinematic viscosity (Table 1). In this setting, at steady state the Nusselt number is unity and the average internal temperature at steady state in an internally heated convective fluid (Parmentier and Sotin, 2000) follows a power scaling law

$$T/T_H = c Ra_H^{-\beta}, \quad (4)$$

with c and β dimensionless constants, and $T_H = \rho H R^2 / k$ the internal heating temperature scale with R now the fluid thickness in the experiments. Experiments yield $\beta = 1/4$ at steady state (Limare et al., 2015), in agreement with theory (Vilella and Kaminski, 2017). In principle, this law holds for the temperature difference across the unstable boundary layer. It turns out, however, that the temperature at the edge of the boundary layer and the average internal temperature can both be scaled to $Ra_H^{-1/4}$ and hence are proportional to one another (Vilella et al., 2018; Limare et al., 2015). We confirmed that this scaling is valid in our experiments over several orders of magnitude in Ra_H as can be seen in the accompanying MethodsX paper (Limare et al., 2020) with $c = 3.45 \pm 0.15$.

As we consider the transient regime in the experiments, i.e., before the steady state is reached, a secular term dT/dt has to be introduced in the scaling law. Assuming that the secular term can be treated as an additional contribution to internal heating, equation (4) becomes

$$T = c Ra_H^{*- \beta} \rho \left(H - C_p \frac{dT}{dt} \right) R^2 / k, \quad (5)$$

where Ra_H^* is a modified Rayleigh-Roberts number defined as

$$Ra_H^* = \frac{\rho g \alpha (H - C_p \frac{dT}{dt}) R^5}{k \kappa \nu}. \quad (6)$$

The two previous equations can be combined to yield an expression for the time derivative of the average internal temperature,

$$\rho C_p \frac{dT}{dt} = \rho H - \left(\frac{kT}{cR^2} \right)^{\frac{1}{1-\beta}} \left(\frac{g \alpha R^5}{k \kappa \nu} \right)^{\frac{\beta}{1-\beta}}. \quad (7)$$

Further noting that the conservation of energy per unit surface writes

$$\rho C_p R \frac{dT}{dt} = \rho H R - Q_s, \quad (8)$$

we obtain an expression for the surface heat flux in the convective regime,

$$Q_s = \left(\frac{kT}{cR} \right)^{\frac{1}{1-\beta}} \left(\frac{g \alpha R^4}{k \kappa \nu} \right)^{\frac{\beta}{1-\beta}}, \quad (9)$$

that is found in very good agreement with the experimental results over several orders of magnitude (Limare et al., 2020). We will now apply these experimentally validated theoretical results to systems where the secular evolution of the average internal temperature is also related to the time evolution of internal heating rate as in the case of a planetesimal.

3.1. Adaptation of the scaling-laws to the thermal evolution of a planetesimal

To apply the experimentally determined scaling law to the modeling of the thermal evolution of a planetesimal, we need to consider several differences between the experimental setting and the natural case. First, we multiply the literal expression we have obtained in equation (9) by a factor 1/3 corresponding to a spherical geometry, as established by Deschamps et al. (2012). Second, we take into account the decreasing heating rate induced by the progressive disintegration of ^{26}Al and the implications on the material properties of a planetesimal given in Table 1.

In the parent body, viscosity is not constant but changes as a function of the average internal temperature and – when relevant – the fraction of melt, and is given by

$$\mu = \mu_0 f(\phi^{\text{melt}}) \exp\left(\frac{E_{\text{act}}}{R_g T} - \frac{E_{\text{act}}}{R_g T_0}\right), \quad (10)$$

where μ_0 is a reference viscosity at $T_0 = 1000^\circ\text{C}$ (that will be set between 10^{18} and 10^{21} Pa.s, see section 4), R_g is the ideal gas constant, E_{act} is the activation energy (Table 1), and $f(\phi^{\text{melt}})$ is defined as

$$f(\phi^{\text{melt}}) = \exp(-\sigma_{\text{sil}} \phi_{\text{sil}}^{\text{melt}}) \exp(-\sigma_{\text{Fe}} \phi_{\text{Fe}}^{\text{melt}}), \quad (11)$$

where the fractions of melt are taken as a linear function of temperature,

$$\phi_{\text{sil,Fe}}^{\text{melt}} = \frac{T - T_s^{\text{sil,Fe}}}{T_l^{\text{sil,Fe}} - T_s^{\text{sil,Fe}}}, \quad (12)$$

if $T_s^{\text{sil,Fe}} < T < T_l^{\text{sil,Fe}}$ with $T_s^{\text{sil,Fe}}$ and $T_l^{\text{sil,Fe}}$ the solidus and liquidus temperature of silicate and metal, respectively, and $\sigma_{\text{Fe}} = 4$ and $\sigma_{\text{sil}} = 21$ (Scott and Kohlstedt, 2006; Hustoft et al., 2007).

Because in silicate bodies the viscosity is strongly temperature dependent, we must also account for the presence of a conductive viscous lid at the top of the convective system (e.g. Davaille

and Jaupart, 1993). In that case, four additional parameters have to be introduced: the thickness of the lid and the temperature drop across the lid, the thickness of the thermal boundary layer at the top of the convective fluid and the temperature drop across the thermal boundary layer. The thickness and temperature drop across the thermal boundary layer at the top of the convective fluid are given by well established scaling laws for thermal convection with large viscosity variations (Davaille and Jaupart, 1993; Choblet and Sotin, 2000). These laws introduce a “viscous scale” ΔT_{visc} for the temperature,

$$\Delta T_{\text{visc}}(T) = -2.21 \frac{\mu(T)}{d\mu/dT(T)}. \quad (13)$$

The thickness δ_{lid} of the viscous lid and the temperature T_{lid} at its base are then obtained by imposing the continuity of temperature and heat flux at the base of the lid. To that aim we use $T = T_{\text{lid}} + \Delta T_{\text{visc}}$ as the average internal temperature of the convective fluid and $\delta_{\text{lid}} = kT_{\text{lid}}/Q_s$, while replacing R by $R - \delta_{\text{lid}}$ in equation (9).

The energy absorbed by partial melting, E_{lat} , and the energy brought by surface impacts during accretion, E_a , are the last ingredients to take into account in the conservation of energy in the case of a planetesimal. The rate of heat absorption by partial melting is given by

$$E_{\text{lat}} = L_{\text{sil}} \frac{d\xi_{\text{sil}}}{dt} + L_{\text{Fe}} \frac{d\xi_{\text{Fe}}}{dt}, \quad (14)$$

with L_{sil} and L_{Fe} the latent heat of the silicate and metal (Table 1), respectively, and ξ_{sil} and ξ_{Fe} their melt fraction. The rate of variation of the melt fractions for $T_s^{\text{sil,Fe}} < T < T_l^{\text{sil,Fe}}$ is derived from equation (12) and writes

$$\frac{d\xi_{\text{sil,Fe}}}{dt} = \frac{dT/dt}{T_l^{\text{sil,Fe}} - T_s^{\text{sil,Fe}}}, \quad (15)$$

with $T_s^{\text{sil}} = 1150^\circ\text{C}$ and $T_s^{\text{Fe}} = 990^\circ\text{C}$ the silicate solidus and the Fe-Fe eutectic, respectively, and $T_l^{\text{sil}} = 1800^\circ\text{C}$ and $T_l^{\text{Fe}} = 1615^\circ\text{C}$ for the silicate liquidus and the pure Fe melting temperature, respectively. To take into account surface impacts, we further consider that an impactor of mass M_i and temperature T_i brings a gravitational energy $E_G = GMM_i/R$, with M and R the mass and radius of the target, respectively, and a thermal energy $E_T = M_i C_p T_i$. The fraction of energy that is transferred to the target rather than radiated into space, E_a , is then set to a nominal fraction of 30% of $(E_G + E_T)$ similar to the value used by Sramek et al. (2012). The effect of this fraction is discussed in the Supplementary Material Fig. 11.

4. Accretion curve of the parent bodies of the magmatic iron meteorites from their thermal evolution

4.1. Instantaneous accretion

As a first step in the study of the thermal evolution of planetesimals we consider, as in previous models, instantaneous accretion and a purely conductive model. To be consistent with the sizes inferred from the metallographic cooling rates we calculate the thermal evolution of a 300 km body as a function of its time of accretion. We further consider a constant temperature of 0°C at the surface of the planetesimal. The results shown in the inset of Fig. 2 imply that the parent bodies of non-carbonaceous magmatic iron meteorites accreted earlier (≈ 0.5 Myr after CAIs formation) than the ones of carbonaceous magmatic iron meteorites (≈ 1 Myr after CAIs formation). These results are consistent with the conclusions obtained for a 40 km body by Kruijer et al. (2014) which shows that within the framework of instantaneous accretion and a purely




Fig. 2. Evolution of the average internal temperature of a 300 km radius planetesimal as a function of its time of instantaneous accretion, and for two values of the reference viscosity μ_0 . The red circles give the position of the “non-carbonaceous” magmatic iron meteorites and the blue circles of the “carbonaceous” magmatic iron meteorites. Compared to a purely conductive model (dashed lines in the inset, with colors corresponding to the same age of formation as in the main figure) the average internal temperature is much lower when convection is taken into account. This is due to more effective heat transfer to the surface in the convective regime than in the conductive one.

conductive model the size of the body is not a key parameter for the thermal evolution.

Conductive models predict extreme temperatures in the planetesimals, larger than the liquidus of the metal and silicate phases, hence total melting and differentiation of the non-carbonaceous magmatic iron meteorites (inset in Fig. 2). Hence at least melting-induced convection should be taken into account in the thermal modeling. Both the increase of temperature and the occurrence of melting actually decrease the viscosity (Supplementary Material Fig. 8) and increase the value of the Rayleigh-Roberts number in the planetesimal. We obtain that the critical value for convection, $Ra_{H,cr} = 5758$ for a rigid boundary condition at the surface (Schubert et al., 2001), is reached at ≈ 0.1 Myr after CAIs (if accretion also starts at CAIs). Once $Ra_H \geq Ra_{H,cr}$ convection should start. However, in a fluid with a strongly temperature dependent viscosity the onset of convection is significantly delayed (Davaille and Jaupart, 1993; Choblet and Sotin, 2000). We calculate the onset time of convection using the scaling given by experiments and numerical simulations (Davaille and Jaupart, 1993; Choblet and Sotin, 2000),

$$t_0 = \frac{1}{\pi} \frac{R^2}{\kappa} \left(\frac{Ra_{H,cr}}{Ra_H} \right)^{-2/3} \left(\frac{T}{\Delta T_{visc}} \right)^{8/3}. \quad (16)$$

The onset time evolves as a function of time following the increase of temperature in the planetesimal. We consider that convection starts when the age of the body becomes larger than t_0 and we obtained an onset time of ≈ 0.2 Myr. This can be taken as an upper bound as convection may start as soon as some melt segregation of iron triggers a density contrast driven Rayleigh-Taylor instability.

Thermal evolution in the case of instantaneous accretion, when convection is taken into account, is shown in Fig. 2. Convection increases heat transfer in the body and results in significantly cooler average internal temperatures than in the purely conductive case. But the main conclusions reached in the conductive case still hold: the parent body of non-carbonaceous magmatic iron meteorites accreted earlier and underwent a hotter thermal evolution leading to total melting and complete differentiation. Convection does




Fig. 3. Model results for continuous pebble accretion. The red circles give the position of the “non-carbonaceous” magmatic iron meteorites and the blue circles of the “carbonaceous” magmatic iron meteorites (Kruijer et al., 2014; Budde et al., 2016; Yang et al., 2008, 2010); time is given by the Hf-W age. The accretion curves are a function of the initial (nucleus) radius of the parent body. The green and magenta correspond to a nucleus size of 35 km and 90 km, respectively. The dash-dotted lines give the purely conductive solution, while the dashed and solid lines give the complete solutions with a convective stage for a reference viscosity μ_0 of 10^{19} and 10^{18} Pa s, respectively. The stars indicate the onset on convection for a reference viscosity of 10^{18} Pa s. For nucleus radii smaller than ≈ 35 km the thermal evolution implies a too large sintering of the parent body and is not consistent with our model framework.

not rule out the capability of instantaneous accretion models to explain the magmatic iron meteorites recorded ages and temperatures. Instantaneous accretion cannot however yield the common thermal evolution we have proposed Fig. 1 and we will look now for accretion scenarios that can produce it.

4.2. Continuous pebbles accretion

Using the time evolution of the average internal temperature proposed in Fig. 1, we invert the energy balance equation (2) as a function of the thermal regime (conduction or convection) to obtain the accretion curves $R(t)$. In this case $E_{int} = E_{lat}$ since E_a can be considered 0 for pebble accretion. Within our framework, the free parameter in the inversion routine is the initial size of the planetesimal, R_0 , that we will refer to as the nucleus, produced by gravitational collapse at the early beginning of the accretion. Because this phenomenon is very rapid compared to the time scale of the thermal evolution recorded by the magmatic iron meteorites, it can be considered as instantaneous in the present model.

In the first stage of accretion, the radius of the body is small and heat transfer occurs by conduction. For the sake of the argument we first consider a purely conductive regime to obtain $R(t)$ from the required thermal evolution (Fig. 1). The results of the inversion are shown in Fig. 3 for a nucleus radius of 35 and 90 km. Based on these results we can calculate the evolution of Ra_H during the accretion of the body (Supplementary Material Fig. 9) and further consider the influence of convection on the model predictions. We calculate the onset time of convection along the accretion curve using the scaling of equation (16). We consider that convection starts (hence that Equation (9) applies) when the age of the body becomes larger than the onset time.

Fig. 3 shows two examples of accretion curves resulting from the inversion of the time evolution of the average internal temperature once convection has been introduced. The shape of the accretion curve is the same as in the case of a purely conductive regime (notably because the initial thermal regime is always




Fig. 4. Universal dimensionless accretion curve predicted by the model (colored thick curve) compared to the parameterized accretion law (dashed white line) of the form $(R(t)/R_\infty)^3 = 1 - (1 - [R_0/R_\infty]^3) \exp(-[t/t_a]^b)$, where $t_a = 1.39 \pm 0.15$ Myr is the timescale of accretion, $b = 1.45 \pm 0.15$ is a shape parameter, and R_0 and R_∞ the initial and final radius, respectively, with $R_0/R_\infty = 0.37 \pm 0.04$. The color of the curves gives the average internal temperature reached in the parent body at a given time (or equivalently for a given radius). The thickness of the colored curve corresponds to the second order variation of the accretion history for different sizes of the initial nucleus.

conductive). However, because convection corresponds to a more efficient heat transfer, it is not possible in this regime to reach the thermal plateau observed at large times in Fig. 1 if cold pebbles continue to accrete at the surface of the planetesimal.

In some cases it is even necessary that the radius of the body decreases in order to counterbalance convection-induced cooling. This is due to the fact that the internal energy varies with volume (which depends on the third power of radius) and the cooling rate is proportional to the surface (second power of radius). Due to convection the body might cool too fast, therefore the model calculates a radius that start to diminish in order to restore the almost constant temperature. This effect is more important for low values of initial radius (cooling/internal heat $\sim 1/R$).

This decrease of radius can correspond either to some erosion, or to sintering at the end of accretion due to compaction of the planetesimal following partial melting. In the following we shall reject the accretion curves yielding to more than 10% of sintering as this is the bulk porosity value we consider in the present model. Based on all the constraints we conclude that the nucleus size of the parent body was between 35 and 90 km (Fig. 3).

As already mentioned, accretion curves depend only on the initial (nucleus) size $R(t=0) = R_0$. When made dimensionless by dividing the radius of the body by its final value, all acceptable solutions (i.e., fitting both the thermal evolution recorded by the magmatic iron meteorites and their sizes) appear to share the same evolution (Fig. 4). This implies that both the shape of the accretion curve, and the associated thermal evolution, do not depend on the final size of the body (which itself depends on the initial nucleus radius), and the growth curves in Figs. 3 follow a single parameterized accretion law (Fig. 4). Three stages of evolution can be introduced to describe the universal dimensionless accretion curve: (i) an initial quasi-instantaneous accretion of a nucleus with $35 \leq R_0 \leq 90$ km, too fast to be resolved at the scale of our model, (ii) a following slower accretion that reaches a plateau or

even ends at ≈ 3 Myr, and (iii) a sintering due to the compaction of the planetesimal following partial melting. Compared to the more simple case of a purely conductive regime, the end of accretion at ≈ 3 Myr appears well resolved in the complete model including convection.

5. Discussion

5.1. Alternative accretion scenarios

To evaluate further the robustness of our model predictions, we study two alternative accretion scenarios: (i) successive collisions with impactors with sizes of tens of kilometers, and (ii) the fragmentation of a parent body.

In scenarios of discontinuous accretion, the planetesimal grow through successive collisions with impactors of a given size R_i . Here we consider that the impactors are formed with their final size R_i at $t = 0$ and we follow their average internal temperature using the conservation of energy as given by equation (2). We model two different types of accretion: orderly or runaway (Fig. 5). The resulting thermal evolution is illustrated Fig. 5 for a final parent body radius of 300 km and $R_i = 40$ km or $R_i = 10$ km. None of these scenarios yield a thermal evolution consistent with the magmatic iron meteorites record.

The results of scenario of discontinuous accretion – in the case of large impactors – are a function of the fraction of energy that is transferred to the parent body during accretion. We study the influence of this parameter for the thermal evolution corresponding to orderly accretion of impactors of 40 km formed at time 0 producing a final body of 300 km in 4 My (Supplementary Material Fig. 11). We obtain that in the case of a perfect energy transfer to the parent body, or in the case of total loss of the impactor energy to space, the thermal evolution predicted is not consistent with the constraints provided by the magmatic iron meteorites. We can then conclude that our preferred scenario of continuous accretion is both robust and the only one able to yield a thermal evolution consistent with the constraints provided by the magmatic iron meteorites.

In the alternative scenario of fragmentation, the planetesimal has a large size at time 0 and is later fragmented into smaller bodies. For the sake of the argument we consider here an initial body size of 300 km, but the conclusions we reach are general. Because of the large initial size, the early thermal evolution is very fast, the surface heat loss ($\propto R^2$) being much smaller than the internal heat sources ($\propto R^3$). As a consequence total melting occurs early, even before the formation of IAB (Supplementary Material Fig. 12). The most favorable end member fragmentation scenario would be a very early fragmentation into small bodies, but this would imply in turn a subsequent episode of continuous accretion and would be equivalent to the formation of a nucleus followed by continuous pebbles accretion.

5.2. Sensibility of the model of pebbles accretion with respect to model parameters

Fig. 3 shows that taking into account convection in a planetesimal under continuous accretion has a small impact on model prediction, limited to an end of accretion delayed from ≈ 3 Myr to ≈ 4 Myr. This reflects the strong control of the model outputs by the thermal evolution of Fig. 1. As discussed above, this rather “cold” thermal evolution results mainly from the accretion of cold pebbles, and the accretion curve is only tuned as a function of the thermal regime. This implies that the model predictions are robust and do not change significantly when we change the parameters used in the scaling laws. For example, a change of the reference viscosity by an order of magnitude has an almost negligible effect




Fig. 5. Alternative accretion scenarios. Upper panel: evolution of the radius of the planetesimal as a function of time for the two types of accretion considered: orderly (blue curve) and runaway (red). Lower panel: corresponding thermal evolution. The thin black line gives the evolution of the temperature of the impactor in the case of orderly accretion. (a, c) Results for 40 km impactors: the initial thermal evolution is too fast and leads to massive melting not consistent with the geochemical constraints. (b, d) Results for 10 km impactors: orderly accretion generates a scenario consistent with the parent bodies of NC iron magnetic meteorites only.

(Fig. 3). We obtain also a negligible effect of a 50% change in the values of activation energy and of melt sensitivities σ_{Fe} and σ_{sil} , or for a surface temperature of -150°C instead of 0°C (Supplementary Material Fig. 10).

We also allowed the system to differentiate, considering iron segregation and migration towards the center of the parent body to form a metallic core. There are three effects of differentiation for the convecting silicate mantle: (i) a prefactor in the Rayleigh number (Equation (3)) equal to $(1 + f + f^2)/3$, where $f = R_{core}/R$ (Deschamps et al., 2012) with R_{core} the core radius given by the mass fraction of metal, (ii) a concentration of the heat source, hence an increase of $H(t)$, in the mantle, as no Aluminium is partitioned into the core, and (iii) a decrease of its thickness from R to $D = R - R_{core}$. The two last effects tend to counterbalance each other and the resulting bulk impact of the differentiation on the accretion curves is not significant, as can be seen in Fig. 6.

5.3. Comparison with astrophysical models

The first two stages of our accretion law are consistent with astrophysical models that predict (i) a fast initial accretion due to

gravitational collapse of meter to kilometer size boulders (Johansen et al., 2015; Simon et al., 2017) and (ii) a rather cold and continuous snowball sweeping of dust and pebbles (Xie et al., 2010). The second stage, corresponding to pebble accretion, can further be interpreted in terms of the evolution of the fraction of pebbles in the disk.

Visser and Ormel (2016) performed numerical simulations of pebble accretion as a function of the sizes of the accreted particles and of the accreting planetesimal as well as of the distance from the star. They define and compute a pebble accretion time scale,

$$t_{growth} = \frac{M}{\dot{M}} = \frac{4\rho_* R}{3v_{hw} \rho_p f_{coll}}, \quad (17)$$

with M the mass of the accreting planetesimal, $\dot{M}$ its time derivative, $\rho_* = 1 \text{ g cm}^{-3}$ the internal density of the pebbles, $v_{hw} = 5780 \text{ cm s}^{-1}$ the magnitude of the headwind faced by the planetesimal, f_{coll} the collisional efficiency, and ρ_p the density of pebbles in the disk (expressed as a relative density compared to the gas density ρ_g). The value of t_{growth} depends on the size of the planetesimal, on the solid-to-gas ratio ρ_p/ρ_g , and on the distance




Fig. 6. Evolution of the accretion curves when differentiation is taken into account (dotted lines) compared to the reference evolution without differentiation (solid lines). In these calculation differentiation occurs instantaneously when the solidus of the silicate is reached ($T = 1150^\circ\text{C}$). Differentiation produces a larger sintering at the end of the accretion in small bodies.

from the star. For the sake of the argument, we shall use the results obtained by Visser and Ormel (2016) at 3 AU and for a typical planetesimal size of 100 km which is relevant for our study and corresponds to a minimal dependence of the result on the size of the pebbles. For this choice of parameters, and for a typical solid-to-gas $\rho_p^{\text{ref}}/\rho_g = 1/100$, Visser and Ormel (2016) show in their figure 4 that the growth timescale $t_{\text{growth}}^{\text{ref}}$ ranges between 40 Myr and 100 Myr for pebbles radii between 0.01 cm and 30 cm. These time scales are one order of magnitude larger than the one estimated by our model. This implies that the solid-to-gas ratio during the accretion of iron meteorites was also one order of magnitude larger.

For a given accretion curve $R(t)$ obtained by the inversion of the thermal evolution recorded by the magmatic iron meteorites it is possible to calculate the evolution of the accretion time scale $t_{\text{growth}}(t) = \frac{1}{3}R(t)/\dot{R}(t)$. In the conditions described above, the ratio $t_{\text{growth}}(t)/t_{\text{growth}}^{\text{ref}}$ is directly the ratio $\rho_p^{\text{ref}}/\rho_p(t)$ with $\rho_p(t)$ the time-varying fraction of pebbles seen by the planetesimal during its accretion. As an example, we show in Fig. 7 the evolution of $\rho_p(t)/\rho_g$ for a nucleus of 65 km. As expected, the density of pebbles starts with a very high value, consistent with the production of a nucleus by gravitational collapse in overdense zones in the disk, and sharply decreases around 3 Myr at the end of the accretion. One should note however that this marks also the appearance of a new mechanism (sintering and/or erosion) that would artificially decrease the effective density of pebbles “seen” by the planetesimal.

5.4. Consequences for the differentiation of the parent body of magmatic iron meteorites

Under canonic nebular conditions, iron can be anticipated to be present within these pebbles and dust as metallic Fe (dominant form), FeS and oxidized Fe in silicates (Ebel, 2006). Upon heating within the planetesimal, it is expected that FeS melts first at the corresponding Fe-FeS eutectic $\approx 1000^\circ\text{C}$ (Brett and Bell, 1969) (see Supplementary Material, Fig. 13). As the temperature rises, the melt becomes more iron rich. As Kruijer et al. (2014) we further assume that the metallic melt remains in contact with the silicate matrix until the total melting of FeS alloy. Under this assumption,




Fig. 7. Evolution of the density of pebbles in the disk at 3 AU for a nucleus of 65 km that grows by pebble accretion following the accretion curve based on the inversion of the thermal record of magmatic iron meteorites for a reference viscosity $\mu_0 = 10^{19}$ Pa s. Two values of the pebbles radius r_p are considered, 0.01 cm and 30 cm. The pebbles density is very high at the beginning of the accretion – in agreement with the hypothesis of boulders accretion in dense parts of the disk – and sharply decreases to values $< 10^{-2}$ by the end of accretion at ≈ 3 Myr after CAIs. The two steps in the solid line are due to the sharp decrease of viscosity in the planetesimal when iron and then silicate start to melt.

the recorded temperature corresponds to the liquidus temperature set by the Sulfur content and the Hf-W age to the time at which this liquidus temperature is reached. This implies that the systematic difference in age between the old “non-carbonaceous” (NC) magmatic iron meteorites and the younger “carbonaceous” (CC) magmatic iron meteorites, is not due to a difference in accretion age, but simply related to the fact that NC are richer in S, and thus reached their FeS liquidus temperature earlier than CC.

It is worth noting that metallic melt separation that occurs at liquidus temperature does not necessarily correspond to total differentiation, thus the Hf-W age does not correspond to the age of core formation, but rather to the formation of metal pockets large enough to prevent metal-silicate W re-equilibration. Eutectic melting of Fe-FeS can take place much below the temperature of melting of the silicate framework which may indeed hamper the percolation of the Fe-FeS liquids to the core. The problem of the differentiation of an accreting planetesimal has been previously addressed using a purely conductive model, pure Fe metal only and no surface tension (Sramek et al., 2012) for a 500 km planetesimal that accreted within 3 Myr. Their results can be used to discuss the process of compaction and core formation in our scenario. At 3 My, about half of the metal is molten and has segregated to form a central core and a non negligible part of the molten metal is furthermore trapped in the solid silicate matrix as lenses at intermediate depths. It is very likely that metal extraction will be even less efficient in our model for two reasons. First, all things being equal, the average internal temperature in a convective planetesimal (our case) is smaller than in a conductive one (the case studied in Sramek et al. (2012)). Hence the rate of production of molten metal, and the subordinated rate of metal extraction, will be smaller in the convective than in the conductive regime. As a consequence differentiation will take a longer time in a convective planetesimal and will have reached a less advanced stage when cooling and solidification will occur. Second, it has been theoretically established that high surface tension (such as the one between solid silicate and Fe-FeS metal) enhances local melt concentration rather than global melt extraction (Ricard et al., 2001). Experiments have further shown that a threshold fraction

of about 15 vol% of Fe-S melt was required to overcome surface tension effects and to extract the melt from the solid silicate matrix (Bagdassarov et al., 2009). We can thus conclude that in our model complete melting of the Fe-FeS mixture is a necessary condition for an efficient compaction leading to the formation of a core. However, melting of more than 50% of the silicate matrix will be further necessary to yield complete metal extraction and full differentiation (Sramek et al., 2012), as in that case percolation is not limited by the rheology of the matrix anymore.

Based on the previous arguments, we conclude that the parent bodies of magmatic iron meteorites might be only partially differentiated, some iron meteorites being possibly fragments of local ponds of molten metal rather than pieces of the central core (Fig. 4). This conclusion is in fact not specific to the present accretion model. In the models considered up to now for the formation of the parent bodies of iron meteorites, i.e., instantaneous accretion at given times (Kruijer et al., 2014), the thermal evolution calculated taking into account convection predicts much lower average internal temperatures than previously anticipated in purely conductive models (Fig. 2). In a scenario of instantaneous accretion, total melting and full differentiation occurred only for the “non carbonaceous” magmatic iron meteorites.

The present accretion model, while developed to reproduce the thermal histories of the parent bodies of magmatic iron meteorites, seems to explain also correctly two major characteristics of the parent bodies of CV carbonaceous chondrites, namely (i) the age of the end of their accretion and (ii) the coexistence of an external undifferentiated shell with a central convective metallic core potentially powering a dynamo (Gattacceca et al., 2016). In our scenario, convection develops in the planetesimal underneath a stable conductive viscous lid of a few kilometer thickness (Supplementary Material Fig. 14). The interior of this viscous lid is by definition much colder than the interior of the body hence cannot reach the solidus and remains undifferentiated. The distribution of ^{26}Al ages in chondrules from CV chondrites implies that accretion of their parent body was still ongoing ≈ 4 Myr after CAIs (Villeneuve et al., 2009). In our model this would correspond to the very last stage of limited accretion occurring between ≈ 3 and 4 Myr, once sintering has become the dominant feature. At 4 Myr, the 300 °C isotherm in the viscous lid is located ≈ 2 –3 km below the surface, allowing the preservation of primitive accreted materials including chondrules and of undisturbed ^{26}Al isochrons in these chondrules (Supplementary Material Fig. 14). Such a planetesimal would have developed a metallic core which could remain partially melted long enough for a dynamo to develop in agreement with recent paleomagnetic observations in chondrules from CV chondrites (Gattacceca et al., 2016). It is worthwhile noting that the presence of a stagnant lid and of internal heating reduce the amount of heat that can be extracted from the core and transported by convection to the surface. In some cases, if the internal heating is too high, the heat flux at the boundary between the core and the mantle may even be negative, i.e. the core is heating up, which would then help to keep it molten for longer periods of time.

6. Conclusion

We have developed a model for the transient thermal evolution of a planetesimal in the course of accretion. Compared to previous approaches, the model takes into account both a conductive regime (at the beginning of accretion when temperature increase and radius are small) and a convective regime. Although convection is not necessarily required by our model, for nominal values of the parameters that enter the experimentally derived scaling laws, convection would start around 1 Myr during the accretion.

We show that the thermal evolution recorded by all the magmatic iron meteorites can be described by a unique exponential law that we latter employed to obtain accretion curves using the conservation of energy in the parent bodies of the magmatic iron meteorites. Once made dimensionless, all the accretion curves define a universal accretion law that introduces three stages: (i) instantaneous nucleus formation at $t = 0$ by gravitational collapse, (ii) continuous pebble accretion at a rate decreasing with time and ending at ≈ 3 Myr, (iii) negligible accretion accompanied by a few percent sintering by compaction.

A main difference of our model compared to previous predictions of models that considered instantaneous accretion and a purely conductive thermal regime is a colder thermal evolution controlled by the continuous accretion of cold “pebbles”, and that may yield partial differentiation only. Some of the magmatic iron meteorites may thus be local metal ponds rather than the remnant of the metallic core of their parent body.

Furthermore, regardless the cooling mode (conduction or convection) in the parent body, there will always exist a cold viscous stagnant lid of a few kilometers that should have remained completely undifferentiated.

On another hand the presence of a stagnant lid may also delay the cooling of the planetesimal even if cooling proceeds by convection. If such a planetesimal is differentiated, then the metallic core can develop a dynamo for a non-negligible amount of time.

Partial melting of the silicate phase will produce a certain degree of silicate differentiation in addition to the formation of an iron core. Hence multi-phase flow modeling will be required to further describe the thermal evolution of the planetesimal during that stage. Such a complex modeling remains however beyond the scope of this study.

More generally, our formalism which combines convective thermal transfer and the evolution of the size of the body during its formation, could be used to compute the thermal evolution produced by a given accretion law, hence to provide additional constraints for theoretical models of planetesimals formation and evolution.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

We would like to thank the reviewers and the editor William B. McKinnon for their fruitful comments that improved the manuscript. This work was funded by the ANR-11-IS04-0004 project and supported by the Tellus/PNP program of INSU-CNRS. Many thanks to Medi@Terre for improving Fig. 4. This project contributes to the IdEx project Université de Paris ANR-18-IDEX-0001.

Appendix A. Supplementary material

Supplementary material related to this article can be found online at <https://doi.org/10.1016/j.epsl.2020.116469>.

References

- Bagdassarov, N., Golabek, G.J., Solferino, G., Schmidt, M.W., 2009. Constraints on the Fe-S melt connectivity in mantle silicates from electrical impedance measurements. *Phys. Earth Planet. Inter.* 177, 139–146.
- Birnstiel, T., Fang, M., Johansen, A., 2016. Dust evolution and the formation of planetesimals. *Space Sci. Rev.* 205, 41–75. <https://doi.org/10.1007/s11214-016-0256-1>.
- Brett, R., Bell, P.M., 1969. Melting relations in the Fe-rich portion of the system Fe-FeS at 30 kb pressure. *Earth Planet. Sci. Lett.* 6, 479–482.

- Budde, G., Burkhardt, C., Brennecke, G.A., Fischer-Gödde, M., Kruijer, T.S., Kleine, T., 2016. Molybdenum isotopic evidence for the origin of chondrules and a distinct genetic heritage of carbonaceous and non-carbonaceous meteorites. *Earth Planet. Sci. Lett.* 454, 293–303.
- Carlsaw, H.S., Jaeger, J.C., 1959. *Conduction of Heat in Solids*. Oxford University Press, New York.
- Choblet, G., Sotin, C., 2000. 3D thermal convection with variable viscosity: can transient cooling be described by a quasi-static scaling law? *Phys. Earth Planet. Inter.* 119, 321–336.
- Cuzzi, J.N., Hogan, R.C., Shariff, K., 2008. Toward planetesimals: dense chondrules clumps in the protoplanetary nebula. *Astrophys. J.* 687, 1432–1447.
- Davaille, A., Jaupart, C., 1993. Transient high-Rayleigh-number thermal convection with large viscosity variations. *J. Fluid Mech.* 253, 141–166.
- Delbo, M., Walsh, K., Avdellidou, C., Morbidelli, A., 2017. A major asymmetric dust trap in a transition disk. *Science* 357, 1026–1029. <https://doi.org/10.1126/science.aam6036>.
- Deschamps, F., Yao, C., Tackley, P.J., Sanchez-Valle, C., 2012. High Rayleigh number thermal convection in volumetrically heated spherical shells. *J. Geophys. Res.* 117, E09006.
- Ebel, D.S., 2006. Condensation of rocky materials in astrophysical environments. In: Lauretta, D.S., McSween, H.Y.J. (Eds.), *Meteorites and the Early Solar System II*. University of Arizona Press, Tucson, pp. 253–277.
- Gattacceca, J., Weiss, B.P., Gounelle, M., 2016. New constraints on the magnetic history of the cv parent body and the solar nebula from the Kaba meteorite. *Earth Planet. Sci. Lett.* 455, 166–175.
- Ghosh, A., Weidenschilling, S.J., McSween Jr, H.Y., 2003. Importance of the accretion process in asteroid thermal evolution: 6 Hebe as an example. *Meteorit. Planet. Sci.* 38, 711–724.
- Goldstein, J.I., Scott, E.R.D., Chabot, N.L., 2009. Iron meteorites: crystallization, thermal history, parent bodies, and origin. *Chem. Erde* 69, 293–325.
- Haack, H., Rasmussen, K.L., Warren, P.H., 1990. Effects of regolith megaregolith insulation on the cooling histories of differentiated asteroids. *J. Geophys. Res.* 95, 5111–5124.
- Hevey, P.J., Sanders, I.S., 2006. A model for planetesimal meltdown by ^{26}Al and its implications for meteorite parent bodies. *Meteorit. Planet. Sci.* 41, 95–106.
- Hustoft, J., Scott, T., Kohlstedt, D.L., 2007. Effect of metallic melt on the viscosity of peridotite. *Earth Planet. Sci. Lett.* 260, 355–360.
- Johansen, A., Lambrechts, M., 2017. Forming planets vis pebble accretion. *Annu. Rev. Earth Planet. Sci.* 12451, 359–387.
- Johansen, A., Mac Low, M.M., Lacerda, P., Bizzarro, M., 2015. Growth of asteroids, planetary embryos, and Kuiper belt objects by chondrule accretion. *Sci. Adv.* 1, e1500109. <https://doi.org/10.1126/sciadv.1500109>.
- Kleine, T., Munker, C., Mezger, K., Palme, H., 2002. Rapid accretion and early core formation on asteroids and the terrestrial planets from Hf-W chronometry. *Nature* 418, 952–955.
- Kruijer, T.S., Burkhardt, C., Budde, G., Kleine, T., 2017. Age of Jupiter inferred from the distinct genetics and formation times of meteorites. *Proc. Natl. Acad. Sci. USA* 114, 6712–6716.
- Kruijer, T.S., Touboul, M., Fischer-Gödde, M., Bermingham, K.R., Walker, R.J., Kleine, T., 2014. Protracted core formation and rapid accretion of protoplanets. *Science* 344, 1150–1154. <https://doi.org/10.3847/2041-8213/aa8c79>.
- Lee, D.C., Halliday, A.N., 1997. Core formation on Mars and differentiated asteroids. *Nature* 388, 854–857.
- Lichtenberg, T., Golabek, G.J., Gerya, T.V., Meyer, M.R., 2016. The effects of short-lived radionuclides and porosity on the early thermo-mechanical evolution of planetesimals. *Icarus* 274, 350–365. <https://doi.org/10.1016/j.icarus.2016.03.004>.
- Limare, A., Kenda, B., Kaminski, E., Surducan, E., Surducan, V., Neamtu, C., 2020. Transient lab-scale experiments of internally heated convection. Submitted to *MethodsX*.
- Limare, A., Vilella, K., Di Giuseppe, E., Farnetani, C.G., Kaminski, E., Surducan, E., Surducan, V., Neamtu, C., Fourel, L., Jaupart, C., 2015. Microwave-heating laboratory experiments for planetary mantle convection. *J. Fluid Mech.* 777, 50–67.
- van der Marel, N., et al., 2013. A major asymmetric dust trap in a transition disk. *Science* 340, 1199–1202.
- McSween, H.Y.J., 1999. *Meteorites and Their Parent Planets*. Cambridge University Press, Cambridge.
- Merk, R., Breuer, D., Spohn, T., 2002. Numerical modeling of ^{26}Al -induced melting of asteroids considering accretion. *Icarus* 159, 183–191. <https://doi.org/10.1006/icar.2002.6872>.
- Morbidelli, A., Raymond, S.N., 2016. Challenges in planet formation. *J. Geophys. Res., Planets* 121, 1962–1980. <https://doi.org/10.1002/2016JE005088>.
- Ormel, C.W., Klahr, H.H., 2010. The effect of gas drag on the growth of protoplanets. analytical expressions for the accretion of small bodies in laminar disks. *Astron. Astrophys.* 520, A43.
- Parmentier, E.M., Sotin, C., 2000. Three-dimensional numerical experiments on thermal convection in a very viscous fluid: implications for the dynamics of a thermal boundary layer at high Rayleigh number. *Phys. Fluids* 12, 609–617.
- Qin, L., Dauphas, N., Wadhwa, M., Masarik, J., Janney, P.E., 2008. Rapid accretion and differentiation of iron meteorite parent bodies inferred from ^{182}Hf - ^{182}W chronometry and thermal modeling. *Earth Planet. Sci. Lett.* 273, 94–104.
- Ricard, Y., Bercovici, D., Schubert, G., 2001. A two-phase model for compaction and damage 2. Applications to compaction, deformation, and the role of interfacial surface tension. *J. Geophys. Res.* 106, 8907–8924.
- Schubert, G., Turcotte, D.L., Olson, P., 2001. *Mantle Convection in the Earth and Planets*. Cambridge University Press, Cambridge.
- Scott, T., Kohlstedt, D.L., 2006. The effect of large melt fraction on the deformation behavior of peridotite. *Earth Planet. Sci. Lett.* 216, 177–187.
- Simon, J.P., Armitage, P.J., Youdin, A.N., Rixin, L., 2017. Evidence for universality in the initial planetesimal mass function. *Astrophys. J. Lett.* 847, L12. <https://doi.org/10.3847/2041-8213/aa8c79>.
- Sramek, O., Millesi, L., Richard, Y., Labrosse, L., 2012. Thermal evolution and differentiation of planetesimals and planetary embryos. *Icarus* 217, 339–354.
- Suzuki, A., Ohtani, E., Kato, T., 1998. Density and thermal expansion of a peridotite melt at high pressure. *Phys. Earth Planet. Inter.* 107, 53–61.
- Vilella, K., Kaminski, E., 2017. Fully determined scaling laws for volumetrically heated convective systems, a tool for assessing habitability of exoplanets. *Phys. Earth Planet. Inter.* 266, 18–28.
- Vilella, K., Limare, A., Jaupart, C., Farnetani, C., Fourel, L., Kaminski, E., 2018. Fundamentals of laminar free convection in internally heated fluids at values of the Rayleigh–Roberts number up to 10^9 . *J. Fluid Mech.* 846, 966–998.
- Villeneuve, J., Chaussidon, M., Libourel, G., 2009. Homogeneous distribution of Al-26 in the solar system from the Mg isotopic composition of chondrules. *Science* 325, 985–988.
- Visser, R.G., Ormel, C.W., 2016. On the growth of pebble-accreting planetesimals. *Astron. Astrophys.* 586 (A66), 1–11.
- Weidenschilling, S.J., 1977. Aerodynamics of solid bodies in the solar nebula. *Mon. Not. R. Astron. Soc.* 180, 57–70.
- Xie, J.W., Payne, M.J., Thébaud, P., Zhou, J.L., Ge, J., 2010. From dust to planetesimals: the snowball phase? *Astrophys. J.* 724, 1153–1164.
- Yang, J., Goldstein, J.I., 2006. Metallographic cooling rates of the IIIAB iron meteorites. *Geochim. Cosmochim. Acta* 70, 3197–3215.
- Yang, J., Goldstein, J.I., Michael, J.R., Kotula, P.G., Scott, E.R.D., 2010. Thermal history and origin of the IVB iron meteorites and their parent body. *Geochim. Cosmochim. Acta* 74, 4493–4506.
- Yang, J., Goldstein, J.I., Scott, E.R.D., 2008. Metallographic cooling rates and origin of IVA iron meteorites. *Geochim. Cosmochim. Acta* 72, 3043–3061.